

ST KYROLLOS FAMILY CLINIC

DR. ASHRAF SADDIK
PN 65028KB
MB.B.Ch,
Diploma Anaesthesia,
FRACGP.
Master Psychopharmacology.
Advanced Diploma CBT.



2A Moore Street, Coburg VIC 3058
Tel (03) 9386 0900 Fax (03) 9386 5388
ASHRAF SADDIK P/L ACN 060 815 401
ABN 20 060 815 401

www.stkyrollosclinic.com.au
www.circumcisiondoctor.com.au
drsaddik@stkyrollosclinic.com.au
drsaddik@circumcisiondoctor.com.au
argus-skfc@netspace.net.au (Encrypted E-mail)

After the circumcision, the first few weeks of the healing process:-

After the circumcision has healed, it is usually necessary to separate the shaft skin from the head of the penis. If that is not done, smegma can accumulate under the adherent skin.

Therefore, the patient needs to return for review after the circumcision has healed.

Check for fresh bleeding, and if this occurs in the first 24 hours, please get in touch with the clinic immediately. Bleeding after the first 24 hours is usually only a drop or two and of no consequence. However, the penis usually does get red and swollen, which is part of the healing process. The circumcised penis will have two layers of dressings.

The bandages will stay longer for a circumcised penis, which is no problem. If the dressing has not dropped off after a day or two days, The **Circumcision doctor**, Dr. Saddik, may remove them if you have trouble removing them.

Leave the dressing around the shaft of the penis for 24 hours and use a sponge to clean the baby. After 24 hours, give the baby a warm bath with salt and gently remove the gauze. After removing the dressing, the circumcised penis is gently washed with soft, warm salt water.

The Circumcision Doctor, Dr. Saddik, is always available to review your baby as often as you feel necessary. If you have any worries, don't hesitate to contact us.

For about one (1) week after the procedure, you will need to place a dab of Vaseline at the tip of your baby's penis at each nappy change. Other than that, treat your baby as usual. It is essential to bring him back after the circumcision has healed (about four weeks after the procedure). Don't neglect this. Use pain relief medication such as paracetamol for infants, only if needed. If the wound bleeds during the first 24 hours after the procedure, apply pressure with an ice pack in a clean cloth for 20 minutes and contact the clinic on the number shown in "Contact us."